

Marine Dissolved Organic Phosphorus Composition: Insights from Samples Recovered  
Using Combined Electrodialysis/Reverse Osmosis

A Thesis  
Presented to  
The Academic Faculty

by

Cindy Jackson

In Partial Fulfillment  
of the Requirements for the Degree  
Master of Science in the  
School of Earth and Atmospheric Sciences

Georgia Institute of Technology  
August 2009

Marine Dissolved Organic Phosphorus Composition: Insights from Samples Recovered  
Using Combined Electrodialysis/Reverse Osmosis

Approved by:

Dr. Ellery Ingall, Advisor  
School of Earth and Atmospheric Sciences  
*Georgia Institute of Technology*

Dr. Irina Sokolik  
School of Earth and Atmospheric Sciences  
*Georgia Institute of Technology*

Dr. Josef Dufek  
School of Earth and Atmospheric Sciences  
*Georgia Institute of Technology*

Date Approved: June 26, 2009

For Papaw

## **ACKNOWLEDGEMENTS**

I would like to thank my family and friends for all their love and support. I also wish to thank my advisor and other members of the EAS faculty for providing me with direction and encouragement. Special thanks go to my cats, who are eight warm, fuzzy balls of stress relief. Finally, I would like to thank God for always being so good to me.

# TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iv
LIST OF TABLES	vi
LIST OF FIGURES	vii
LIST OF SYMBOLS AND ABBREVIATIONS	viii
SUMMARY	ix
<u>CHAPTER</u>	
1 INTRODUCTION	1
2 METHODS	3
Sampling locations	3
Sample collection	3
Electrodialysis/reverse osmosis technique	4
<sup>31</sup> P NMR	5
3 RESULTS AND DISCUSSION	8
Polyphosphate	10
Phosphate Esters	14
Phosphonates	17
4 CONCLUSIONS	20
REFERENCES	21

## LIST OF TABLES

	Page
Table 1: Location, depth, recovery and relative abundance of P compound classes	8

## LIST OF FIGURES

	Page
Figure 1: $^{31}\text{P}$ NMR chemical shifts	6

## LIST OF SYMBOLS AND ABBREVIATIONS

DOM	Dissolved Organic Matter
ED/RO	Electrodialysis/reverse osmosis
UF	Ultrafiltration
HMW	High Molecular Weight
LMW	Low Molecular Weight
$^{31}\text{P}$ NMR	Phosphorus Nuclear Magnetic Resonance
DOP	Dissolved Organic Phosphorus
TDP	Total Dissolved Phosphorus
ATP	Adenosine Triphosphate
GTP	Guanosine Triphosphate
DIP	Dissolved Inorganic Phosphorus



## SUMMARY

The dominant organic phosphorus compound classes were characterized in marine samples using a new, high recovery method for isolating and concentrating bulk dissolved organic matter (DOM) called combined electrodialysis+reverse osmosis (ED/RO). In contrast to earlier studies which use ultrafiltration (UF) to recover only the high molecular weight DOM, ED/RO is capable of isolating both low molecular weight (LMW) and high molecular weight (HMW) DOM. Samples were collected from a broad range of marine environments: along a transect incorporating coastal and offshore waters off the Southeastern United States, in Effingham Inlet, a Pacific fjord located on Vancouver Island, British Columbia and in the Amundsen Sea, Antarctica. Results from phosphorus nuclear magnetic resonance ( $^{31}\text{P}$  NMR) analysis reveals a similar abundance of P compound classes between samples, phosphate esters (80-85%), phosphonates (5-10%) and polyphosphates (8-13%). These samples contain significantly higher proportions of polyphosphate P and P esters and lower proportions of phosphonates than measured in previous studies using the UF method. The much higher levels of polyphosphate detected in our samples suggests that polyphosphate is present mainly in the LMW DOM fraction. Polyphosphates in DOM may be present as (or derived from) dissolved nucleotides or organismal polyphosphate bodies, or both. Low molecular weight P esters are likely composed of phosphoamino acids and small carbohydrates, like simple sugar phosphates and/or dissolved nucleotides. Phosphonates in DOM are more prevalent as HMW phosphonate compounds, which suggests that LMW phosphonates are more readily utilized in marine ecosystems. Overall, the investigation of DOM across a

size spectrum that includes both the HMW and the LMW fractions reveals a new picture of phosphorus distribution, cycling and bioavailability.

# CHAPTER 1

## INTRODUCTION

Phosphorus is an essential nutrient for life, and the availability of phosphorus in ocean surface waters is an important control on ocean productivity. In surface waters of vast oligotrophic ocean regions, dissolved organic phosphorus (DOP) typically comprises the major fraction (~75-80%) of the total dissolved P (TDP) pool (Karl and Björkman 2002; Benitez-Nelson 2000; Karl and Yanagi 1997; Smith et al. 1986; Jackson and Williams 1985; Orrett and Karl 1987; Bjorkman and Karl 1994). Consequently, regeneration of P from DOP has been shown to play a significant role in supplying the P necessary for biological production (e.g. Karl and Bjorkman 2001; Dyhrman et al. 2006; Karl et al. 2008). Many P studies have focused on identifying specific individual biomolecules, such as adenosine triphosphate (ATP), that typically comprise a small percentage of total P in the ocean (Karl and Bjorkman 2002; Benitez-Nelson 2000). Other studies have investigated the bulk distribution of different P compound classes (e.g. phosphonates, P esters) in samples covering a specific molecular weight range that represent about 25% of the total P in the ocean (Clark et al. 1998, 1999; Kolowitz et al. 2001; Sannigrahi et al. 2006). Because these studies have focused on relatively small fractions of the DOP pool, the bulk composition of marine DOP remains largely unknown.

Uncertainty in DOP composition is a major impediment in our attempts to understand not only the global P cycle but also the coupled cycles of many other elements associated with marine production. DOM plays a number of central roles in

oceanic biogeochemical cycles. It acts as both a conduit and a storehouse for nutrients and reduced carbon. Yet despite a great deal of research conducted over the past two decades into DOM composition and cycling, many questions remain unanswered. Most of these questions, such as the identities and sources of labile LMW dissolved molecules interacting with bacterial and photoautotroph populations, have remained difficult to examine because we lacked the methods necessary to concentrate such ephemeral compounds up to the levels at which they can be examined by modern analytical techniques.

However, a new technique called combined electrodialysis+reverse osmosis (ED/RO) recovers two to three times more DOM from seawater than earlier techniques, thereby allowing us to investigate DOM across a much larger molecular size spectrum than previously possible (Vetter et al. 2007; Gurtler et al. 2008; Koprivnjak et al. 2006). We demonstrate in this paper that the P composition of this more complete DOM isolate is quite different from that observed in previous studies of UF-extracted DOM, and it provides exciting new possibilities for rewriting our understanding of DOM sources, dynamics, and interactions with other organic and inorganic pools.

## CHAPTER 2

### METHODS

#### 2.1 Sampling locations

With our small-scale prototype ED/RO equipment, sample processing can be quite lengthy. Therefore, we were only able to process a limited number of samples for this study. In order to gain a preliminary view of the potential range in marine P composition, samples were collected from five sites in four very different marine environments, ranging from oligotrophic Gulf Stream waters to the eutrophic coastal environments of Georgia and Vancouver Island to highly productive waters of the Antarctic Amundsen Sea. The Gulf Stream sampling site is located off the coast of Savannah, Georgia. The Gulf Stream is a powerful current on the western boundary of the North Atlantic subtropical gyre and is characterized by warm, oligotrophic waters. Coastal samples were taken from Effingham Inlet and Savannah, Georgia at Skidaway Institute of Oceanography. Effingham Inlet is a Pacific fjord, located on the west coast of Vancouver Island in Barkley Sound.

#### 2.2 Sample collection

All samples were collected from a Niskin bottle rosette sampler. A 200 L Gulf Stream sample was collected aboard the *R/V Savannah* in July 2008. Effingham Inlet samples were taken in April and July 2007 aboard the *R/V Barnes*. Two large volume samples of 50 L and 120 L were collected from Effingham Inlet at depths of 61 m and 78 m, respectively. Similarly, two large samples of 50 L and 200 L were collected just off

the coast of Savannah, Georgia. A 200 L Antarctic sample was collected in the Amundsen Sea in December 2008 aboard the *Icebreaker Oden*.

### 2.3 Electrodialysis/reverse osmosis technique

Coupled ED/RO extraction of DOM from seawater samples was performed generally following the protocols described in Vetter et al. (2007). Briefly, all samples and rinse solutions were passed through a 0.45  $\mu\text{m}$  filter. The system was cleaned for the Effingham samples by circulating a minimum 50 L of sample through both the ED and RO systems for 1 hour. For all other samples at least two, 1-hour long, 50 L system rinses were performed. ED is used to remove ions from the seawater sample. In ED, the seawater sample (to be deionized) is contacted with a concentrate flow (to receive ions) across many pairs of cation and anion exchange membranes. A DC electrical current is directed through the membrane stack. Only ionized dissolved components are targeted by ED. The ion exchange membranes are non-porous above the size of the interstitial spaces between the polymer chains in the cross-linked hydrated ion exchange resin. The process results in a concentrate solution, which is discarded, and a deionized solution (diluate). The diluate solution is subsequently concentrated via RO in order to isolate dissolved molecules for analysis.

Concentrated, desalted samples were transferred into clean 10-L HDPE bottles and frozen immediately. Both the RO and ED systems were rinsed with 0.01 M NaOH to recover adsorbed fractions of the DOM in a sample. System mass balances were calculated to determine DOM recoveries using the protocols described in Koprivnjak et al. (2009). The current applied to the system was constantly monitored to make sure that

it remains below the limiting current density determined for the specific system geometry and flow characteristics (Vetter et al. 2007).


#### 2.4 $^{31}\text{P}$ NMR

After samples were freeze dried and ground, bulk P composition was determined with the use of a solid-state  $^{31}\text{P}$  NMR instrument at the NMR center in the School of Chemistry and Biochemistry at the Georgia Institute of Technology. The instrument is a Bruker DSX 400 spectrophotometer and uses Cross Polarization- Magic Angle Spinning (CP-MAS) at a  $^{31}\text{P}$  frequency of 161 MHz. About 90 mg of powdered sample was packed into 4 mm diameter cylindrical zirconia rotor fitted with Kel-F cap and spun at 10,000 +/-10 Hz in a Bruker magic-angle spinning probe (Sannigrahi and Ingall 2005).

For all samples, a cross-polarization sequence, optimized to obtain semiquantitative data, was used with a 1.0 ms contact time and a pulse delay of 4 s (Sannigrahi et al. 2005). For each sample, at least 8000 transients were collected. After the spectra were obtained, peak areas were integrated off-line using the Spinworks software package. Areas were determined by dropping vertical lines to the baseline in a similar fashion as that of Hedges et al. (2002). Gaussian fits and other more complicated spectral deconvolution techniques were not employed because little information is available on shapes and positions of peaks in natural samples, which is needed in order to account for contributions from each overlapping P bonding environment.

Functional groups within the sample were recognized by comparing chemical shift values relative to an external phosphoric acid standard. Chemical shift values are dimensionless and are given in parts per million (ppm) relative to the phosphoric acid standard, which is set at 0 ppm. These values are essentially variations in resonance

frequency of different compound classes compared to a standard. To determine the percentage of sample belonging to each phosphorus functional group, the spectra were integrated over the following chemical shifts in ppm (Figure 1): -15 to -25 (polyphosphate), 7 to -15 (phosphate esters), 30 to 15 (phosphonate), 69 to 47 and -55 to -77 (spinning side bands).



**Figure 1** The  $^{31}\text{P}$  NMR chemical shifts corresponding to different P compound classes, superimposed on the  $^{31}\text{P}$  NMR spectrum of an Effingham Inlet sample taken at 78m. Asterisks denote spinning side bands



Spinning side bands are an artifact of magic-angle spinning and are added to the appropriate associated peak area. The chemical shift range used to determine polyphosphate in this paper is conservative compared to other papers which report it between -18 and -30 ppm (e.g. Sannigrahi and Ingall 2005). The analytical error of  $^{31}\text{P}$  NMR as determined from previous studies is +/-10% of the measured value (Sannigrahi and Ingall 2005; Hedges et al. 2002; Sannigrahi et al. 2005).

## CHAPTER 3

### RESULTS AND DISCUSSION

A comparison of the relative abundances of three P compound classes (polyphosphates, P esters, and phosphonates) between samples collected with ED/RO and samples collected with UF in previous studies yield three key observations (Table 1): 1) Polyphosphates are not readily observed in samples collected using UF but are observed in all samples collected by ED/RO. 2) P esters comprise the largest fraction of the DOP pool for both collection techniques. 3) Phosphonates make up a smaller portion of DOP in ED/RO samples as compared to UF samples.

**Table 1** Location, depth, recovery and relative abundance of P compound classes for each site

\* Kolowith et al. 2001; Sannigrahi et al. 2006; Clark et al. 1999

Sample	Location	Sampling Depth (m)	% Recovery	% P ester	% phosphonate	% polyphosphate
Skidaway Dock	31°59' N, 81°1'W	1	61	85	7	8
Effingham Inlet	48°55'N, 125°2'W	78	77	80	7	13
Effingham Inlet	48°55'N, 125°2'W	61	68	85	5	10
Gulf Stream	30°48N, 80°19'W	5	84	80	9	11
Amundsen Sea	72°46'S 115°24' W	250	54	82	10	8
UF Literature*	Many regions	All Depths	~25	~75	~25	0

The dominance of P esters in the DOM pool is consistent with previous investigations of the HMW fraction recovered by UF, however differences in the relative abundance of other P compounds such as phosphonates and polyphosphates to P esters between ED/RO and UF samples are undoubtedly related in part to the different molecular size fractions recovered by the two techniques.

Although differences abound between samples recovered using ED/RO compared to UF samples from the literature in similar locations, perhaps the most intriguing finding is the rather limited compositional variation in ED/RO samples between the vastly different sites sampled so far. This finding is in close agreement with the results of Kolowitz et al. (2001), in which no variation was found in the relative proportions of P compound classes in HMW DOM recovered from several oligotrophic ocean regions and at all depths. Furthermore, our results show striking similarities between very oceanographically different regions – eutrophic and oligotrophic. These observations seem to indicate that DOP is relatively the same compositionally throughout the world ocean.

It is perplexing that regions with varying types of organisms and nutrient levels would exhibit invariance in abundance of P compound classes, but upon closer examination we can reconcile these findings with current knowledge of marine P cycling. P is cycled rapidly in marine environments (Smith et al. 1985; Karl and Bjorkman 2002; Benitez-Nelson 2000; Benitez-Nelson and Buesseler 1999). Thus, labile forms of P are quickly removed, leaving a pool of less labile material with a slow turnover time (Clark et al. 1998, 1999; Kolowitz et al. 2001; Karl and Bjorkman 2002). Although each environment contains different organisms, similar molecules must be preferentially

selected for decomposition and produced during decomposition processes, which leads to the P distribution observed.

### **3.1 Polyphosphate**

Polyphosphates have been identified in marine particulate organic matter (Paytan et al. 2003), but polyphosphates have never before been recognized in NMR spectra of marine DOM. Ultrafiltration samples contain only the HMW dissolved polyphosphate fraction, but samples recovered using ED/RO contains both HMW and LMW fractions. Because polyphosphate can exist in a range of molecular sizes depending on the length of the linear polyphosphate polymer, we expected to find dissolved polyphosphates in a continuum of molecular weights. However, polyphosphates were only present in the LMW. For this reason, we suspect that the amount of HMW polyphosphate recovered by conventional UF techniques is not zero as previously reported, but that it is too low to be detected. Comparison of UF samples to ED/RO samples indicates that polyphosphates are most prevalent in the LMW fractions of DOM. The presence of polyphosphates in LMW DOM is surprising especially in nutrient-poor open ocean environments, such as the Gulf Stream, where such P-rich molecules would be a valuable potential resource for microorganisms. The presence of LMW polyphosphates in such environments suggests LMW polyphosphates are in forms that are difficult for organisms to utilize.

A polyphosphate compound consists of three or more orthophosphate groups linked together in a chain by phosphoanhydride (P-O) bonds. According to the current understanding of polyphosphates in cells, polyphosphates can be produced by organisms via mechanisms involving luxury uptake, or the uptake of P in excess of an organism's immediate need. Luxury uptake allows organisms to form polyphosphates when

another nutrient other than P limits growth, thereby permitting the organism to accumulate excess P reserves that would become highly valuable under future potential P starvation. One mechanism by which organisms can store excess P is by constructing chains of phosphate, which are encapsulated into granules for later use. Biogenic polyphosphate granules are complexed with charge-balancing cations, principally divalent cations such as calcium, and are stored within the cell cytoplasm, until the organism requires it. When the cell lyses, the polyphosphate granules can be released back into the water column, where the granules may remain stable until they are used by other organisms.

Polyphosphate can be found in two forms inside a typical cell: as storage granules and as nucleotides such as adenosine triphosphate (ATP). The larger-sized granules should be recovered by both UF and ED/RO. However, the largest percentage of dissolved polyphosphate in seawater appears within the LMW portion, and this LMW portion can only be recovered by ED/RO. While these LMW polyphosphates may be primarily comprised of small (<1000 Daltons) storage granules, the nucleotide interpretation of LMW dissolved polyphosphate is also consistent with the molecular nature of these compounds. For example, ATP weighs about 500 Daltons. The cutoff for UF is 1000 Daltons, making nucleotides impossible to recover unless ED/RO is used.

Given the rapid cycling of LMW polyphosphates in numerous cellular biochemical reactions, it might be presumed that nucleotides should be highly bioreactive and therefore unstable in the water column. However, Bjorkman and Karl (2005) were able to measure concentrations of ATP and GTP (guanosine triphosphate) at Station Aloha in the North Pacific subtropical gyre, near Hawaii, and found these nucleotides to decrease

in concentration with depth. Therefore, despite the fact that the nucleotides are reactive, they are allowed to build up to some quantifiable extent. To this extent, perhaps the utilization of dissolved nucleotides by pelagic organisms is uptake limited.

If we assume typical surface DOP concentrations of 0.1-0.2  $\mu\text{M}$  (Karl and Bjorkman 2002), and use our average polyphosphate content of 11%, a rough calculation indicates that polyphosphate concentrations should be in the nano-molar range. The dissolved nucleotide concentrations measured by Bjorkman and Karl (2005) are on the order of pico-molar, which are too small to account for the dissolved polyphosphate we see. To reconcile the data, it is crucial to understand what is actually being detected by the methods used. The firefly bioluminescence assay employed by Bjorkman and Karl (2005) is only able to detect specific nucleotides, but NMR is sensitive to all tri-nucleotides. Any molecule with three or more phosphate groups in a chain (not only ATP and GTP) is designated a polyphosphate by NMR. One possibility is that LMW polyphosphate is derived from degraded nucleotides, which may be less bioavailable than the original nucleotides and can no longer be characterized as such.

Furthermore, it has long been hypothesized that certain mechanisms can transform organic phosphorus molecules into inorganic ones. Photochemical and enzymatic processes could possibly break the ester bond, cleaving off the base of the nucleotide. Organisms in need of an organic substrate may take up the base, leaving only the triphosphate chain behind. Photochemical reactions are potentially important in the first 100 meters or so, where light is able to penetrate. Enzymes such as alkaline phosphatase and nucleotidases are responsible for breaking apart the ester bonds in nucleotides. Alkaline phosphatase is produced by all marine organisms, and laboratory studies show

that marine bacteria begin to produce this enzyme at DIP concentrations much higher than typical surface ocean DIP concentrations (Hassan and Pratt 1977). Therefore, although alkaline phosphatase is less efficient than nucleotidases at the specific task of breaking ester bonds in nucleotides, it is ubiquitous in world oceans and may be a key player in the alteration of organic compounds. Thus, attribution of the polyphosphate signals to nucleotides and degradation products of nucleotides is as feasible as the granule explanation for the source of LMW polyphosphate in DOM.

Ultimately, the oceanographic importance of LMW dissolved polyphosphate depends on how easily organisms may utilize it. Although we can hypothesize that the structure of this polyphosphate is much like that of ATP, which is readily used by organisms, the most important control on bioavailability may be the size of the molecules. For example, ATP weighs about 500 Daltons and is small enough to be taken up directly into the cell. Once inside, ATPases can break down the phosphoanhydride bonds within the ATP molecule in order to utilize phosphorus, or the molecule may be kept intact to be used in metabolic processes. However, every organism has a limit on the size of molecule that it can take up directly. For example, Weiss et al. (1991) found that the outer membrane of gram negative bacteria can only transport molecules up to about 600 Daltons. Therefore, if we consider granules as a source, it is possible that these polyphosphates could be too large to cross cell membranes without first being broken up by extracellular enzymes. This would imply that the bioavailability of these compounds depends on the prevalence of organisms which can manufacture the appropriate enzymes. The uncertainty of the structure and size of dissolved

polyphosphate leaves us with a perplexing lack of knowledge about the nature of these compounds which future research will be directed towards answering.

### **3.2 Phosphate Esters**

In all ED/RO samples, we observe greater amounts of phosphate esters compared to samples recovered by UF (Table 1). UF only recovers molecules from the HMW (> 1000 Daltons) fraction of DOM, while ED/RO isolates both HMW and LMW compounds. Comparison of UF samples to ED/RO samples suggests the LMW fraction is enriched in P esters. LMW P ester molecules, such as glucose phosphate and fructose phosphate, are important to key universal cellular metabolic processes, such as glycolysis. Determining the types of molecules that likely compose these LMW P esters can help provide insights as to the reactivity and oceanographic significance of this compound class.

Phosphate esters are synthesized by all living organisms and are incorporated into numerous common biomolecules such as phosphoamino acids, phosphosugars and P lipids. For this reason, it is not surprising that P esters comprise the most abundant compound class in marine DOP. Because most of the organic matter in the ocean is released by phytoplankton, the dominance of P esters in DOM recovered by ED/RO is undoubtedly related to the abundance of these compounds in plankton cells.

An organism will typically prefer to directly take up dissolved inorganic phosphorus (DIP) from seawater, rather than use a DOP molecule. This is because the use of DOP requires the organism to manufacture an enzyme to break the phosphate out of the organic compound before it can be utilized. All marine organisms share the ability to produce enzymes, such as alkaline phosphatase, which are capable of breaking the C-O-P bond of a phosphate ester (Karl and Bjorkman 2002). Laboratory studies have



shown, however, that marine bacteria begin to produce alkaline phosphatase even when there are already high levels of DIP in the water column (Hassan and Pratt 1977). The ubiquity of alkaline phosphatase and other ester hydrolyzing enzymes in ocean waters leads to large fluxes of DIP derived from ester molecules into the water column.

Upon examining possible associations of biomolecules with P in HMW DOM samples, Sannigrahi et al. (2006) found a strong positive correlation ( $R^2 > 0.80$ ) between P content and amino acids and carbohydrates but no correlation between P and lipid contents. If similar trends extend to the LMW fraction, a correlation between amino acids and carbohydrates might also be expected. Indeed, the addition of phosphate groups to amino acids by a P ester linkage forms phosphoamino acids, which are very small molecules that play an important role in the regulation of cellular metabolism. In bacteria, the amino acids histidine, aspartic acid, and glutamic acid are usually favored for phosphorylation (Sannigrahi et al. 2006; Yan et al. 1998).

It is also likely that many of these LMW P esters are associated with carbohydrates. Carbohydrates comprise 10-70% of the plankton cell (Romankevich 1984) and are largest fraction of organic matter in ocean, accounting for 20-30% in surface waters (Pakulski and Benner 1994). Studies have shown that carbohydrates account for up to 80-90% of total extracellular release by phytoplankton (Myklestad 1995) and that most carbohydrates in the water column come from cells (Biersmith and Benner 1998; Romankevich 1984). Therefore, phytoplankton provides a sizable flux of carbohydrates to the water column.

Of these carbohydrates, it is probable that LMW aldoses and nucleotides constitute a significant portion of LMW P esters. Aldoses play an important role in cellular function

and are likely the bulk of LMW carbohydrates in cells. A study revealed that the largest percentage of aldoses (68%) is found in the LMW fraction of DOM (Skoog and Benner 1997). Because glucose is the dominant aldose in phytoplankton cellular material (Biersmith and Benner 1998), glucose phosphate is likely to make up a large portion of LMW P esters. In fact Skoog and Benner (1997) found glucose to be the most abundant aldose in LMW DOM. However, the same study reports only about 7-20% of carbohydrates have been identified as aldoses, when measuring aldoses using HPAE-PAD (high performance anion-exchange chromatography with pulsed amperometric detection) separation/detection process and total carbohydrate using MBTH (Skoog and Benner 1997). This may be due to the limitations of methods used to measure aldoses in seawater which arise when the compound has been altered. If an aldose has been degraded, it may not be measured by the HPAE-PAD separation/detection technique, but may still be recognized as an aldose by NMR or MBTH if it still contains the functional groups.

Nucleotides contain one P ester bond and a three-member phosphate chain. If the LMW polyphosphate compounds we measured are mainly composed of nucleotides which still have a P ester group attached, another contributor to LMW P ester compounds is nucleotides. The presence of nucleotides in a sample will lead to corresponding increases in the relative abundances of both polyphosphates and P esters, as compared to abundances of UF samples from the literature. This is indeed what we observe, and may be evidence that the LMW P ester abundance has a large contribution from nucleotides.

Many LMW P ester compounds are small enough to be readily utilized and go directly across the cell membrane. Just as the presence of nucleotides in DOM is

enigmatic, so is the existence of these small P esters. The extracellular enzyme, alkaline phosphatase, is needed to utilize these compounds and is ubiquitous in seawater. Therefore, LMW P esters would be expected to be generally reactive. However, they are present in abundance in DOM. Phytoplankton cells provide a large flux to the water column, which may allow the compounds to build up to some extent. An alternate explanation for the presence of LMW P esters is that they are altered to a form that few organisms can utilize, yet may contain the functional groups detectable by NMR.

### 3.3 Phosphonates

When comparing our data to the two previous studies (Table 1), phosphonate abundance in samples processed using ED/RO is about half of that in samples processed by UF. Lower phosphonate abundances in ED/RO samples indicate that 1) LMW phosphonates are not as common as HMW phosphonates in marine organisms; or 2) the distribution reflects the end result of cycling P processes. If the distribution reflects cycling processes, then LMW phosphonates are removed from DOM more readily than HMW phosphonates.

Phosphonates are a group of organic molecules characterized by a direct C-P bond, rather than the more prevalent C-O-P ester linkage found in most phosphorus-containing biomolecules. There are not many direct studies of the occurrence of specific phosphonate compounds in marine settings. The occurrence as detected by NMR is widespread as indicated by the results presented as well as in several studies (Kolowitz et al. 2001; Sannigrahi et al. 2006; Clark et al. 1999). Phosphonate compounds are ubiquitous in marine organisms but typically comprise a very small percentage of the total phosphorus. In fact, biochemical pathways that cycle phosphonates in the ocean are

highly unusual given the low prevalence of phosphonates in marine systems and may be artifacts of ancient metabolic processes (Pasek 2008). Most organisms do not produce the enzymes required to break the C-P bond, resulting in a build-up of phosphonates in the environment. However, research has shown that nitrogen fixing bacteria, such as *trichodesmium*, living in oligotrophic waters have the ability to utilize phosphonates (Dyhrman et al. 2006). Bacteria may also represent an important source of phosphonates in marine systems (Kolowitz et al. 2001).

The specific biomolecules and structures containing phosphonates in marine organisms are not as well defined as polyphosphates and P esters. It has been speculated that at least some of the P in HMW fractions is in the form of phosphonolipids. These phosphonolipids may increase the resistance to degradation of cellular membranes (Kennedy and Thompson 1970; Rosenberg 1973). This chemical resistance may manifest itself in the relative preservation and abundance of these compounds in the HMW P pool. On the other hand, Sannigrahi et al. (2006) did not find a correlation between lipid and P contents in DOM, which may rule out the phosphonolipids as a key contributor to the phosphonates.

It is likely that LMW phosphonate compounds may be equally or more common in cells than HMW phosphonate compounds. Thus, decrease in phosphonate abundance in the ED/RO material relative to UF recovered DOM suggests that LMW phosphonate compounds may be more bioavailable than HMW phosphonate. There have been few direct studies in marine environments, but studies in soils and wastewater treatment systems reveal the utilization of several LMW phosphonate compounds, such as aminoethylphosphonate and methylphosphonate, in cells (Cook et al. 1978; Daughton et

al. 1979). If many of the LMW compounds are small enough to travel across cell membranes, their utilization may in fact be more energetically favorable (Rittenberg and Hespell 1975). Indeed, LMW phosphonate compounds, such as methylphosphonate, are bioavailable to certain organisms, and utilization of these compounds is important on a global level (Karl et al. 2008). Thus, the distribution of phosphonates in DOM recovered by ED/RO likely reflects the aggregate of a number of P cycling processes.

## CHAPTER 4

### CONCLUSIONS

Use of the ED/RO technique has yielded new insights into marine DOP composition and cycling. Samples recovered using ED/RO and analyzed by  $^{31}\text{P}$  NMR reveal similar relative proportions of P esters (80-85%), phosphonates (5-10%) and polyphosphates (8-13%) in marine settings ranging from oligotrophic to eutrophic. We hypothesize these striking similarities between different sites are a result of similar decomposition processes operating in all marine ecosystems, causing the accumulation of similar, less bioavailable P compounds. Of particular interest is the LMW pool, which past methods of isolating DOM have been inadequate at recovering. Virtually all polyphosphates and a sizable portion of P esters in DOM occur in this LMW fraction. Ultimately, the oceanographic importance of the LMW pool will depend on the bioavailability of the mixture of LMW DOP compounds that compose it. Future research will be necessary to better define the distribution and bioavailability of LMW DOP compounds in the ocean.

## REFERENCES

- Benitez-Nelson CR (2000) The biogeochemical cycling of phosphorus in marine systems. *Earth-Sci Rev* 51: 109-135
- Benitez-Nelson CR, Buesseler KO (1999) Variability of inorganic and organic phosphorus turnover rates in the coastal ocean. *Nat* 6727: 502-505
- Biersmith A, Benner R (1998) Carbohydrates in phytoplankton and freshly produced dissolved organic matter. *Mar Chem* 63(1-2): 131-144
- Bjorkman K, Karl DM (1994) Bioavailability of inorganic and organic phosphorus compounds to natural assemblages of microorganisms in Hawaiian coastal waters. *Mar Ecol Prog Ser* 111: 265-273.
- Bjorkman K, Karl DM (2005) Presence of dissolved nucleotides in the North Pacific Subtropical Gyre and their role in cycling of dissolved organic phosphorus. *Aquat Microb Ecol* 39: 193-203
- Clark LL, Ingall ED, Benner R (1999) Marine organic phosphorus cycling: novel insights from nuclear magnetic resonance. *Am J Sci* 2999: 724-737
- Cook AM, Daughton CG, Alexander M (1978) Phosphate Utilization by Bacteria. *J Bacteriol* 133(1): 85-90
- Daughton CG, Cook AM, and Alexander M (1979) Bacterial Conversion of Alkylphosphonates to Natural Products via Carbon-Phosphorus Bond Cleavage. *J Agric Food Chem* 27(6)
- Dyhrman ST, Chappell PD, Haley ST, Moffet JW, Orchard ED, Waterbury JB, Webb EA (2006) Phosphonate Utilization by the globally important marine diazotroph *Trichodesmium*. *Nat* 439: 68-71
- Gurtler B K, Vetter TA, Perdue EM, Ingall E, Koprivnjak JK, Pfromm PH (2008) Combining reverse osmosis and pulsed electrical current electro dialysis for improved recovery of dissolved organic matter from seawater. *J Membr Sci* 323: 328-336
- Hassan HM, Pratt D (1977) Biochemical and Physiological Properties of Alkaline-Phosphatases in 5 Isolates of Marine-Bacteria. *J Bacteriol* 129: 1607-1612
- Hedges JI, Baldock JA, Gelinas Y, Lee C, Peterson ML, Wakeham SG (2002) The biochemical and elemental compositions of marine plankton: A NMR perspective. *Mar Chem* 78: 47-63

- Jackson GA, Williams PM (1985) Importance of dissolved organic nitrogen and phosphorus to biological nutrient cycling. *Deep Sea Res* 32: 223-235
- Karl DM, Beversdorf L, Bjorkman K, Church M, Martinez A, DeLong EF (2008) Aerobic Production of Methane in the Sea. *Nat Geosci* 1(7): 473-478
- Karl DM, Bjorkman K (2001) Phosphorus cycle in seawater: Dissolved and particulate pool inventories and selected phosphorus fluxes. *Methods Microbiol* 30: 239-270
- Karl DM, Bjorkman K (2002) Dynamics of DOP. In: *Biochemistry of Marine Dissolved Organic Matter*, Elsevier Science, USA, pp 249-366
- Karl DM, Yanagi K (1997) Partial characterization of the dissolved organic phosphorus pool in the oligotrophic North Pacific Ocean. *Limnol Oceanogr* 42: 1398-1405
- Kennedy KE, GA Thompson (1970) Phosphonolipids – Localization in Surface Membranes of Tetrahymena. *Sci* 168(3934): 989-991
- Kolowitz LC, Ingall ED, Benner R (2001) Composition and cycling of marine organic phosphorus. *Limnol Oceanogr* 46(2): 309-320
- Orrett K, Karl DM (1987) Dissolved organic phosphorus production in surface seawater. *Limnol Oceanogr* 32: 383-395
- Koprivnjak JF, Perdue EM, Pfromm PH, (2006) Coupling reverse osmosis with electro dialysis to isolate natural organic matter from fresh waters. *Water Res* 40(19): 3385-3392
- Koprivnjak JF, Pfromm PH, Ingall E, Vetter TA, Schmitt-Kopplin P, Hertkorn N, Frommberger M, Knicker H, Perdue EM (2009) Chemical and spectroscopic characterization of marine dissolved organic matter isolated using coupled reverse osmosis–electrodialysis. *Geochim Cosmochim Acta* 73:4215-4231
- Myklestad SM (1995) Release of Extracellular Products by Phytoplankton with Special Emphasis on Polysaccharides. *Sci Total Environ* 165(1-3): 155-164
- Orrett K, Karl DM (1987) Dissolved organic phosphorus production in surface seawater. *Limnol Oceanogr* 32: 383-395
- Pakulski JD and Benner R (1994) Abundance and Distribution of Carbohydrates in the Ocean. *Limnol Oceanogr* 39(4): 930-940
- Pasek MA (2008) Rethinking early Earth phosphorus geochemistry. *Proceedings of The National Academy of Sciences Of The United States of America* 105: 853-858



- Paytan A, Cade-Menun BJ, McLaughlin K, Faul KL (2003) Selective phosphorus regeneration of sinking marine particles: evidence from  $^{31}\text{P}$ -NMR. *Mar Chem* 82: 55-70
- Rittenberg SC, Hespell RB (1975) Energy efficiency of intraperiplasmic growth of *Bdellovibrio bacteriovorus*. *J Bacteriol* 121: 1158-1165
- Romankevich EA (1984) *Geochemistry of Organic Matter in the Ocean*. Springer Verlag, New York, pp 334
- Rosenburg H (1973) Phosphonolipids, In: Ansell GB, Hawthorne JN, Dawson RMC (ed.), *Form and function of phosphonolipids*, Elsevier, Amsterdam, pp 333-344
- Sannigrahi P, Ingall ED, Benner R (2005) Cycling of dissolved and particulate organic matter at station Aloha: Insights from  $^{13}\text{C}$  NMR spectroscopy coupled with elemental, isotopic, and molecular analyses. *Deep Sea Res I* 52(8): 1429-1444
- Sannigrahi P, Ingall E (2005) Polyphosphate as a source of enhanced P fluxes in marine sediments overlain by anoxic waters: Evidence from  $^{31}\text{P}$  NMR, *GeochemTrans*, 6(3): 52-59.
- Sannigrahi P, Ingall E, Benner R (2006) Nature and dynamics of phosphorus-containing components of marine dissolved and particulate organic matter. *Geochim Cosmochim Acta* 70(23): 5868-5882
- Skoog A and Benner R (1997) Aldoses in various size fractions of marine organic matter: Implications for carbon cycling. *Limnol Oceanogr* 42(8): 1803-1813
- Smith REH, Harrison WG, Harris L (1985) Phosphorus exchange in marine microplankton communities near Hawaii. *Mar Biol* 86: 75-74
- Smith SV, Kimmerer J, Walsh TW (1986) Vertical flux and biochemical turnover regulate nutrient limitation of net organic production in the North Pacific Gyre. *Limnol Oceanogr* 31: 161-167
- Vetter TA, Perdue EM, Ingall E, Koprivnjak JF, Pfromm PH (2007) Combining reverse osmosis and electrodialysis for more complete recovery of dissolved organic matter from seawater. *Sep Purif Technol* 56: 383-387
- Wakeham SG, Lee C, Hedges JI, Hernes PJ, Peterson ML (1997) Molecular indicators of diagenetic status in marine organic matter. *Geochim Cosmochim Acta* 61(24): 5363-5369
- Weiss MS, Abele U, Weckesser J, Welte W, Schiltz E, Schultz GE (1991) Molecular architecture and electrostatic properties of a bacterial porin. *Sci* 254: 1627-1630

Yan JX, Packer NH, Gooley AA, William KL, (1998) Protein phosphorylation: technologies for the identification of phosphoamino acids. *J Chromatogr* 80: 23-41